بِسۡمِ ٱللَّهِ ٱلرَّحۡمَٰنِ ٱلرَّحِيمِ

Existence of Dark Matter

By Ayan Ahmad Bhat

November 16, 2025.

Table of Contents

1. Historical Background
2. Key Scientific Statements
3. Theoretical Framework
    - WIMPs
    - Axions
    - Sterile Neutrinos
    - Modified Gravity (MOND)
4. Observational Evidence
    - Galaxy Rotation Curves
    - Cosmic Microwave Background
    - Gravitational Lensing and the Bullet Cluster
5. Detection Methods
    - Direct Detection
    - Indirect Detection
    - Collider Searches
6. Current and Future Research
7. Conclusion and Open Questions
8. References

Abstract

Dark matter remains one of the most profound mysteries in modern astrophysics and cosmology. Although it does not emit, absorb, or reflect light, its presence is revealed through its powerful gravitational influence on galaxies, galaxy clusters, and the large-scale structure of the universe. This thesis presents a comprehensive examination of the historical development of dark matter research, beginning with early observations by Fritz Zwicky and later groundbreaking findings by Vera Rubin. It explores the leading theoretical models—including Weakly Interacting Massive Particles (WIMPs), axions, and sterile neutrinos—as well as alternative theories such as Modified Newtonian Dynamics (MOND). The study further reviews key observational evidence such as galaxy rotation curves, gravitational lensing, cosmic microwave background measurements, and the Bullet Cluster. Current detection methods, including direct, indirect, and collider-based searches, are analyzed to highlight the global scientific effort to uncover the nature of dark matter. Despite decades of research, the true identity of dark matter remains unknown, continuing to inspire new technologies, international collaborations, and future experiments. This thesis underscores the importance of dark matter in shaping our universe and the ongoing pursuit to understand this elusive component of cosmic matter.

Historical Background

The concept of dark matter emerged in the early 20th century. In 1933, Swiss astronomer Fritz Zwicky studied the Coma galaxy cluster and found that the galaxies were moving far too quickly to be bound by the gravity of visible matter alone. He proposed an unseen “dunkleMaterie” (German for *dark matter*) to account for this missing mass. Decades later, in the 1970s, Vera Rubin and colleagues measured the rotation speeds of stars in spiral galaxies and found that stars far from the galactic centers orbit too quickly to be held by luminous matter alone. Rubin “concluded that these galaxies must be held together by dark matter,” and her observations provided such strong evidence that the dark matter hypothesis became widely accepted. By the late 20th century, Zwicky’s and Rubin’s results had convinced the astronomical community that most of the Universe’s mass is invisible.

Following these early discoveries, many other lines of evidence accumulated. Observations of galaxy clusters, gravitational lensing, and cosmological structure consistently required far more mass than could be seen. The term “dark matter” became standard after Zwicky’s coinage, and scientists explored candidates ranging from faint stellar remnants to new particle types. Early suggestions such as ordinary neutrinos were quickly ruled out because known (active) neutrinos are too light and fast to account for the missing mass. By the end of the 20th century, non-luminous matter was firmly recognized as a major component of the cosmos, even though its nature remained mysterious.

Key Scientific Statements

"Ordinary matter makes up only about 5% of the universe, while dark matter makes up about 27%"."Dark matter makes up most of the mass in galaxies and galaxy clusters"."Even though normal matter makes up everything we can see, the universe must contain more than five times as much dark matter to fit the observations"."Cold non-baryonic dark matter appears to make up 85% of the matter … of the universe, and yet we don’t yet know what it is"."WIMPs … are hypothesized to make up most [dark matter] and therefore some 22% of the universe".In galaxy cluster collisions like the Bullet Cluster, “the intracluster gas and dark matter are separated, offering further evidence in support of dark matter”.

These statements highlight that dark matter is by far the dominant form of mass in cosmic structures, yet its composition is unknown. Each quote is drawn from a reputable source (NASA or scholarly review) to underscore the empirical basis of dark matter research.

Theoretical Framework

WIMPs

Weakly Interacting Massive Particles (WIMPs) are one of the leading dark matter candidates. By definition, WIMPs are massive, electrically neutral particles that interact via the weak force and gravity. They are **nonbaryonic** (not made of protons or neutrons) and would have masses at the electroweak scale or above. Many extensions of the Standard Model predict WIMPs: for example, supersymmetry yields neutralinos, and other theories propose heavy particles with names like axions or Kaluza–Klein modes. Because a thermal population of WIMPs in the early universe would naturally freeze out with a relic abundance comparable to the observed dark matter density, this scenario is sometimes called the "WIMP miracle." The key scientific statement that WIMPs "are hypothesized to make up most dark matter" underscores their prominence in theory. Huge experimental efforts (underground detectors, colliders, and telescopes) aim to discover WIMPs by detecting their rare interactions or production.

Axions

Axions are a very different class of hypothetical particles, originally proposed to solve the strong-CP problem in quantum chromodynamics. If they exist, axions would be extremely light (perhaps μeV to meV mass) and interact extremely weakly. Importantly, axions could form cold dark matter through non-thermal production mechanisms (e.g. the misalignment mechanism). As one source explains, "with a mass above 5 μeV … axions could account for dark matter, and thus be both a dark-matter candidate and a solution to the strong CP problem". In practice, this means that axions would behave like a very cold, non-relativistic fluid on cosmic scales. Searches for axions involve resonant cavities (haloscopes) and helioscopes, which look for the tiny conversion of axions to photons in a magnetic field. If axions are the dark matter, experiments like ADMX may eventually detect them.

Sterile Neutrinos

Sterile neutrinos are hypothetical neutral fermions that do not interact via the standard weak force – only gravity (and possibly other unknown forces) – making them "inert" except for their mass. They naturally arise in seesaw models that explain why known (active) neutrinos have very low mass. A sterile neutrino of mass on the order of a few keV could serve as a **warm dark matter** candidate: it would be non-relativistic (cold) by the time of galaxy formation, but could erase small-scale structure below its free-streaming length. In fact, analyses show that if dark matter were sterile neutrinos, they would need masses of order keV to reproduce cosmic structure. Sterile neutrinos could decay into lighter particles (e.g., X-rays) over very long times, so X-ray astronomy is probing this possibility. Thus far, no conclusive signal has been seen, but the concept remains viable.

Modified Gravity (MOND)

As an alternative to particle dark matter, some theories modify the laws of gravity. A famous example is MOND (Modified Newtonian Dynamics), which alters Newton's force law at very low accelerations. MOND was motivated by rotation curve data and can fit many galaxy curves without invoking dark matter. One review notes that "one of the most notable [alternatives] is modified Newtonian dynamics (MOND)". However, modified gravity models generally struggle to fit cluster-scale observations and cosmological data as well as dark matter does. Moreover, evidence like the Bullet Cluster (see below) shows a clear separation between gravitational mass and ordinary mass, favoring particle dark matter over purely modified gravity. In sum, while MOND remains an interesting idea, it is less widely accepted than the dark matter hypothesis.

Observational Evidence

Galaxy Rotation Curves



Figure 1: Observed rotation curve of the Milky Way galaxy (blue) compared to the expected curve (red) from luminous matter alone. In this case, the predicted rotation speed drops off at large radii, but the actual measurements (blue) remain nearly constant. This *flat rotation curve* indicates the presence of additional mass (a dark matter halo) extending beyond the visible disk. Studies consistently show that spiral galaxies have such flat outer curves, implying that "the distribution of mass in the system appears to bear almost no relation to that of light". In other words, there must be substantial unseen mass (dark matter) in the galaxy's halo to hold stars in fast orbits. This discrepancy – between observed velocities and those predicted from visible stars and gas – was one of the first signals of dark matter.

The physics of rotation curves is straightforward. Newtonian gravity would predict that orbital speed $v(r)$ should fall as $\sim r^{-1/2}$ if all mass is interior to the orbit. Instead, most observed spirals have $v(r)$ asymptoting to a constant at large r, indicating an enclosed mass $M(<r) \propto r$. This implies a roughly isothermal halo density $\rho(r)\propto r^{-2}$. The luminous disk contributes only part of the gravitational pull; an extended spherical dark halo is needed to reconcile theory with data. Empirically, Rubin et al. and many successors found that stars at the edges of spirals move "too fast to be held in by the galaxy's luminous matter". Figure 1 illustrates this classic result for the Milky Way: the red (predicted) curve drops off, while the blue (observed) curve remains flat.

Cosmic Microwave Background



Figure 2: All-sky map of the cosmic microwave background (CMB) temperature anisotropy from the WMAP satellite (first-year data). This image shows the tiny fluctuations (at the level of 10^-5) that are the imprint of early-universe sound waves. The angular power spectrum of these anisotropies has several peaks due to acoustic oscillations. Importantly, the relative heights of the first three peaks encode the amounts of normal (baryonic) and dark matter. In particular, precise measurements by WMAP and Planck find that the best-fit cosmological model requires about five times more dark matter than ordinary matter. Equivalently, ordinary baryons are only ~15% of the matter density, with ~85% in a non-luminous form. Thus CMB observations provide strong, quantitative evidence for dark matter consistent with other probes.

Beyond the peak heights, the CMB also constrains dark matter via structure formation. The inferred matter density and its ratio of baryons to dark matter must match both the CMB and later galaxy surveys. The excellent agreement of the Lambda-Cold Dark Matter model (ΛCDM) with CMB data is often cited as one of the clearest pieces of evidence for dark matter. The quoted statement above – "the universe must contain more than five times as much dark matter" to fit WMAP data – quantitatively summarizes this result.

Gravitational Lensing and the Bullet Cluster

In strong gravitational lensing, light from background galaxies is deflected by a massive foreground cluster. This effect depends only on total mass (visible + dark). By mapping the positions and shapes of lensed images, astronomers infer the mass distribution of the cluster. In virtually every cluster studied, the lensing mass greatly exceeds the mass of luminous galaxies and gas. As NASA notes, "all of the visible matter…isn't nearly enough to create the observed warping effects. Dark matter provides the surplus gravity". Lensing thus independently confirms that galaxy clusters have massive dark halos.

A particularly striking case is the **Bullet Cluster** (1E 0657-56), a pair of colliding clusters imaged in X-rays and optical light. Figure 3 shows a Hubble image overlaid with contours of the inferred mass (from lensing). The hot gas (pink X-ray region) experiences drag in the collision and slows down, while the galaxies (blue points) pass through. The lensing mass peaks, however, remain coincident with the galaxies, not the gas, indicating that most of the mass is in a collisionless component. As one study concludes, the Bullet Cluster is “unusual in that the intracluster gas and dark matter are separated, offering further evidence in support of dark matter”. In other words, the unseen dark matter has passed through the collision while interacting only gravitationally. This directly demonstrates that the dominant mass in the clusters is non-luminous and largely non-interacting.

Figure 3: Composite image of the Bullet Cluster. The bulk of the mass (green lensing contours) is aligned with the galaxies, not the pink X-ray emitting gas. This separation is clear evidence for collisionless dark matter.

Gravitational lensing on larger scales (weak lensing surveys) also maps dark matter statistically. For example, cosmic shear measurements of galaxy shapes reveal a web of dark matter filaments. These observations consistently show more clustering than baryons alone could produce, again implying a dominant dark component. In summary, lensing – both strong and weak – provides powerful, direct evidence for dark matter in clusters and large-scale structure.

Detection Methods

Direct Detection

Direct-detection experiments aim to measure dark matter particles scattering off ordinary nuclei in the laboratory. The classic targets are underground cryogenic detectors using heavy nuclei (e.g. xenon, germanium) shielded from cosmic rays. A generic assumption is that dark matter is abundant in the Galactic halo, so the Earth is constantly sweeping through a dark matter wind. Experiments like LUX, XENON1T, and PandaX look for tiny recoil energies from rare WIMP-nucleus collisions. So far, no conclusive signal has been found, but upper limits have reached impressive sensitivities. These searches constrain the WIMP-nucleon cross-section to be below ~10^-47 cm^2 for 100 GeV particles. Over the next decade, larger detectors (XENONnT, LZ, DARWIN) aim to approach the “neutrino floor,” beyond which solar neutrino backgrounds dominate.

Indirect Detection

Indirect-detection searches look for products of dark matter annihilation or decay, such as gamma rays, neutrinos, or antimatter. The idea is that if dark matter (e.g. WIMPs) annihilates in regions of high density (galactic center, dwarf galaxies, galaxy clusters), it will produce excess high-energy particles that telescopes can see. Instruments like the Fermi Gamma-ray Space Telescope, AMS-02, IceCube, and CTA search for unexplained signals. For example, a tentative gamma-ray excess in the Milky Way's center or the 3.5 keV X-ray line from galaxy clusters have been discussed as possible dark matter hints. However, none of these has provided unambiguous confirmation. In general, indirect searches "focus on looking for the products of dark matter interactions (particularly Standard Model particles) rather than the dark matter itself". Non-detection so far places limits on annihilation cross-sections; for instance, Fermi data require WIMP annihilation cross-sections below roughly the thermal relic value for WIMP masses up to ~100 GeV.

Collider Searches

High-energy colliders (notably the LHC) offer a third strategy: try to produce dark matter particles directly. In proton-proton collisions, unknown weakly interacting particles could be created, escaping the detector unseen and carrying away energy. Experiments thus look for events with large missing transverse energy plus a visible recoiling jet or photon. While the LHC has discovered the Higgs boson, it has not yet found any signals of dark-matter candidates (such as supersymmetric particles). Absence of signals sets constraints on many theoretical models. In essence, collider searches attempt to replicate dark matter creation processes that would have occurred in the early universe. Success in this avenue would not only confirm dark matter's existence, but also shed light on its mass and interaction strength.

Current and Future Research

Research on dark matter is intensely active across astrophysics, cosmology, and particle physics. On the astrophysical side, next-generation surveys will map dark matter with unprecedented precision. The Vera C. Rubin Observatory (LSST) and the Euclid mission will map weak gravitational lensing over vast sky areas, improving our knowledge of dark matter's large-scale distribution. NASA's upcoming Nancy Grace Roman Space Telescope will also perform deep weak-lensing surveys and supernova studies, "using so-called weak gravitational lensing that tracks how smaller clumps of dark matter warp the apparent shapes of more distant galaxies". Roman's measurements will probe how structure grows over time, testing whether dark matter is indeed heavy and cold or perhaps lighter and "warm".

On the particle-search side, experiments are upgrading continuously. Direct detectors are scaling up to multi-ton targets to reach the neutrino floor. New technologies (e.g. superheated liquid detectors, quantum sensors) aim to expand the mass range probed. Indirect searches benefit from more data: Fermi continues to observe, CTA (Cherenkov Telescope Array) will soon survey the gamma-ray sky, and space telescopes and X-ray observatories hunt for rare decay signals. The LHC will collect much more data in coming years, and future colliders (like a proposed 100 TeV machine) would dramatically extend the search reach. In theory, novel models are also under exploration, including fuzzy dark matter (ultralight axions forming wave-like structures) and self-interacting dark matter (to solve small-scale structure issues).

In sum, **"dark matter research spans a very broad range of topics and methods"**. While the evidence is overwhelming, the fundamental nature of dark matter remains unknown. The next decade of experiments and observations may finally reveal its identity, or at least narrow down the possibilities substantially.

Conclusion and Open Questions

Dark matter remains one of the central mysteries of modern physics. Observationally, multiple lines of evidence (galaxy dynamics, CMB, lensing, galaxy surveys) all point to a dominant non-luminous component in the Universe. Yet despite its gravitational influence, dark matter has eluded direct detection. The key statements cited above emphasize both how certain we are (e.g. "makes up most of the mass") and how uncertain we remain ("we don't yet know what it is").

Major open questions include: What is the particle (or mechanism) that constitutes dark matter? Is it a single new particle like a WIMP or axion, or a combination (e.g. a "dark sector")? Are there non-gravitational interactions of dark matter at all? Why hasn't any experiment seen it yet, despite sensitive searches? At galaxy scales, puzzles like the "core-cusp" problem and the "too-big-to-fail" problem challenge simple cold-dark-matter predictions. These could hint at new physics (e.g. dark matter self-interactions or feedback processes) or issues in our modeling. There is also the possibility that gravity itself is not fully understood on large scales, though the Bullet Cluster and cosmological data strongly favor dark matter.

In summary, our thesis is that **the preponderance of evidence requires some form of dark matter**, but its nature is undetermined. Addressing this will likely require both new experimental discoveries and deeper theoretical insight. Until then, the dark matter puzzle continues to drive astrophysics and particle physics forward, reflecting fundamental questions about the composition and history of the Universe.

References

- NASA Goddard Space Flight Center. (n.d.). *Dark Matter*. NASA. Retrieved from https://science.nasa.gov/mission/roman-space-telescope/dark-matter/
- NASA Goddard Space Flight Center. (2025, June 30). *NASA Webb 'Pierces' Bullet Cluster, Refines Its Mass*. NASA. Retrieved from https://science.nasa.gov/missions/webb/nasa-webb-pierces-bullet-cluster-refines-its-mass/
- Editors of Encyclopaedia Britannica. (n.d.). *Weakly interacting massive particle (WIMP)*. In *Encyclopædia Britannica*. Retrieved from https://www.britannica.com/science/weakly-interacting-massive-particle
- Green, A. (2025, January 24). *Dark matter: Evidence, theory and constraints*. *CERN Courier*. Retrieved from https://cerncourier.com/a/dark-matter-evidence-theory-and-constraints/
- Wikipedia contributors. (n.d.). *Indirect detection of dark matter*. In *Wikipedia, The Free Encyclopedia*. Retrieved October 2025, from https://en.wikipedia.org/wiki/Indirect_detection_of_dark_matter
- Wikipedia contributors. (n.d.). *Sterile neutrino*. In *Wikipedia, The Free Encyclopedia*. Retrieved October 2025, from https://en.wikipedia.org/wiki/Sterile_neutrino
- Wikipedia contributors. (n.d.). *Axion*. In *Wikipedia, The Free Encyclopedia*. Retrieved October 2025, from https://en.wikipedia.org/wiki/Axion
- Wikipedia contributors. (n.d.). *Galaxy rotation curve*. In *Wikipedia, The Free Encyclopedia*. Retrieved October 2025, from https://en.wikipedia.org/wiki/Galaxy_rotation_curve
- Wikipedia contributors. (n.d.). *Cosmic microwave background*. In *Wikipedia, The Free Encyclopedia*. Retrieved October 2025, from https://en.wikipedia.org/wiki/Cosmic_microwave_background